



Prevention

IMPACT OF AGE ON THE CARDIOVASCULAR EVENT RISK CONFERRED BY HbA1c IN PATIENTS WITH ESTABLISHED CORONARY ARTERY DISEASE

Poster Contributions

Poster Hall B1

Monday, March 16, 2015, 9:45 a.m.-10:30 a.m.

Session Title: Risk Assessment, Exercise and Atrial Fibrillation

Abstract Category: 21. Prevention: Clinical

Presentation Number: 1246-103

Authors: Alexander Vonbank, Christoph Saely, Philipp Rein, Daniela Zanolin, Heinz Drexel, VIVIT Institute, Feldkirch, Austria, Department of Internal Medicine and Cardiology at the Academic Teaching Hospital Feldkirch, Feldkirch, Austria

Background: In the present study we tested the hypothesis that age modulates the impact of HbA1c on cardiovascular event risk in patients with established coronary artery disease (CAD).

Methods: We prospectively recorded cardiovascular events over a mean follow-up period of 4.4 ± 1.2 years in a large consecutive series of 816 patients with angiographically proven CAD, including 376 subjects <65 years and 440 subjects ≥ 65 years.

Results: During follow-up, the incidence of cardiovascular events was 9.3% among subjects <65 years and 24.8% among subjects ≥ 65 years ($p < 0.001$). Among the younger patients, HbA1c strongly and significantly predicted cardiovascular events (HR 1.54 [1.06-2.23]; $p = 0.022$), but not among in the older patients (HR 1.22 [0.94-1.59]; $p = 0.125$). An interaction term age x HbA1c was statistically significant ($p = 0.007$), indicating that HbA1c was a significantly stronger predictor of cardiovascular events among younger than among older CAD patients.

Conclusion: We conclude that HbA1c is a significantly stronger predictor of cardiovascular events in younger patients than in older patients with established CAD.